arXiv:1104.5459v3 [hep-ph] 17 Jan 2013

$K^0_{l3\gamma}$ decays:
branching ratios and T-odd momenta correlations

A.S. Rudenko[1]
Budker Institute of Nuclear Physics
and Novosibirsk State University,
630090, Novosibirsk, Russia

Abstract

The branching ratios of the $K^0 \to \pi^- l^+ \nu_l \gamma$ $(l = e, \mu)$ decays, and the T-odd triple momenta correlations $\xi = \vec{q} \cdot [\vec{p}_l \times \vec{p}_\pi]/M_K^3$, due to the electromagnetic final state interaction, in these processes are calculated. The contributions on the order of ω^{-1} and ω^0 to the corresponding amplitudes are treated exactly. For the branching ratios and T-odd correlation in $K^0 \to \pi^- e^+ \nu_e \gamma$ decay, the corrections on the order of ω are estimated and demonstrated to be small. The results for the branching ratios are in good agreement with the previous ones. The T-odd triple momenta correlations in $K^0_{l3\gamma}$ decays are calculated for the first time. The values of the ξ-odd asymmetry constitute -1×10^{-4} and -4.5×10^{-4} in the $K^0 \to \pi^- \mu^+ \nu_\mu \gamma$ and $K^0 \to \pi^- e^+ \nu_e \gamma$ decays, respectively.

1. The $K^0 \to \pi^- l^+ \nu_l \gamma$ $(l = e, \mu)$ decays were previously studied theoretically in Refs. [1, 2, 3, 4]. Therein the branching ratios of these decays were calculated. As for the T-odd triple momenta correlations $\xi = \vec{q} \cdot [\vec{p}_l \times \vec{p}_\pi]/M_K^3$, as induced by the electromagnetic final state interaction, they were considered only in the $K^+ \to \pi^0 l^+ \nu_l \gamma$ decays [5, 6, 7]; here and below M_K is the kaon mass and $\vec{q}$, $\vec{p}_l$, $\vec{p}_\pi$ are the momenta of γ, l, π, respectively. In principle, these triple correlations can be used to probe new CP-odd effects beyond the Standard Model, which could also contribute to them.

In the theoretical analysis of radiative effects in the discussed processes, the treatment of the accompanying radiation, which gives the effects on the order of ω^{-1} and ω^0 (the last ones originate from the radiation due to the lepton magnetic moment), is straightforward (here and below ω is the photon energy). As to the structure radiation contribution on the order of ω^0, it is also under control, in fact due to the gauge invariance [8]. The contributions on the order of ω (and higher) depend directly on the photon field strength $F_{\mu\nu}$ (and its derivatives) and cannot be fixed in a model-independent way. We assume that the corrections on the order of ω and higher are relatively small. Indeed, more quantitative arguments presented below demonstrate that such contributions into the discussed branching ratios and into the T-odd momenta correlation in $K^0_{e3\gamma}$ decay do not exceed 20%.

2. At the tree level, the $K^0 \to \pi^- l^+ \nu_l \gamma$ decays are described by the Feynman graphs in Fig. 1.

The matrix elements for diagrams 1a and 1b look as follows:

$$M_{1a} = \frac{G}{\sqrt{2}} \sin\theta_c e [f_+(t)(p_K + p_\pi)_\alpha + f_-(t)(p_K - p_\pi)_\alpha] \bar{u}_\nu \gamma^\alpha (1 + \gamma_5) \left(\frac{p_l e^*}{p_l q} + \frac{\hat{q}\hat{e}^*}{2 p_l q} \right) v_l, \tag{1}$$

---

[1] a.s.rudenko@inp.nsk.su



Figure 1: The tree diagrams of $K^0 \to \pi^- l^+ \nu_l \gamma$ decays

$$M_{1b} = -\frac{G}{\sqrt{2}} \sin\theta_c e [f_+(t')(p_K + p_\pi + q)_\alpha + f_-(t')(p_K - p_\pi - q)_\alpha] \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l \frac{p_\pi e^*}{p_\pi q}; \quad (2)$$

here G is the Fermi coupling constant, θ_c is the Cabibbo angle, e is the elementary charge ($e > 0$), $t = (p_K - p_\pi)^2$, $t' = (p_K - p_\pi - q)^2$; here and below the lower indices attached to the matrix elements match the corresponding Feynman diagrams in the figures.

Usually the dependence of the form factors f_+ and f_- on the momentum transfer t is described by formula

$$f_\pm(t) = f_\pm(0) \left(1 + \lambda_\pm \frac{t}{m_\pi^2}\right). \quad (3)$$

The experimental data are adequately described by Eq. (3) with $\lambda_+ \approx 0.03$ for $l = \mu$ and $l = e$; $\lambda_- = 0$ for $l = \mu$ [9]; λ_- for $l = e$ is unknown, but one may assume that it is also close to zero.

In the $K^0_{l3\gamma}$ decays the ratio $\lambda_\pm t/m_\pi^2$ is small, $\lambda_\pm t/m_\pi^2 \lesssim 0.1$, so one can put $f_\pm(t) = f_\pm(0)$. Since the ratio $\xi(0) = f_-(0)/f_+(0) \lesssim 0.1$ is also small [10], one can neglect $f_-(0)$ with the same accuracy.

Thus, our expressions (1) and (2) simplify to

$$M_{1a} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0) (p_K + p_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) \left(\frac{p_l e^*}{p_l q} + \frac{\hat{q}\hat{e}^*}{2 p_l q}\right) v_l, \quad (4)$$

$$M_{1b} = -\frac{G}{\sqrt{2}} \sin\theta_c e f_+(0) (p_K + p_\pi + q)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l \frac{p_\pi e^*}{p_\pi q}. \quad (5)$$

However, the sum of diagrams 1a and 1b is not gauge invariant: it does not vanish under the substitution $e^* \to q$. To restore the gauge invariance, one should add the third diagram where a photon is directly emitted from the vertex (see Fig. 1c). This contact amplitude has no single-particle intermediate states and therefore is on the order of ω^0 and higher. The contribution $\sim \omega^0$, as derived with the Low technique [8], is

$$M_{1c} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0) e^*_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l. \quad (6)$$

Thus, the model-independent gauge invariant tree amplitude of $K^0_{l3\gamma}$ decays, including only terms on the order of ω^{-1} and ω^0 (but all of them!), is

$$M_{tree} = M_{1a} + M_{1b} + M_{1c} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0) \left\{ (p_K + p_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l \left(\frac{p_l e^*}{p_l q} - \frac{p_\pi e^*}{p_\pi q}\right)\right.$$

$$+(p_K+p_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5)\frac{\hat{q}\hat{e}^*}{2p_l q} v_l + \left(e^*_\alpha - \frac{p_\pi e^*}{p_\pi q} q_\alpha\right) \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l \Bigg\} . \quad (7)$$

This expression agrees with the corresponding formulas in Ref. [3] (if our $f_+(0)$ is set to its $SU(3)$ value $f_+(0) = 1$).

It is convenient to present amplitude (7) as a sum of gauge-invariant contributions. They are the "infrared" term M_{IR} corresponding to the sum of the amplitudes of accompanying radiation by the pion and lepton (independent of the lepton magnetic moment), the magnetic term M_{mag} which is the amplitude of spin-dependent accompanying radiation of the lepton magnetic moment, and the Low term M_{Low}:

$$M_{IR} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0)(p_K+p_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l \left(\frac{p_l e^*}{p_l q} - \frac{p_\pi e^*}{p_\pi q}\right), \quad (8)$$

$$M_{mag} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0)(p_K+p_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5)\frac{\hat{q}\hat{e}^*}{2p_l q} v_l, \quad (9)$$

$$M_{Low} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0) \left(e^*_\alpha - \frac{p_\pi e^*}{p_\pi q} q_\alpha\right) \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l. \quad (10)$$

In the lowest order in G the decay width $\Gamma(K^0 \to \pi^- l^+ \nu_l \gamma)$ is equal to the sum $\Gamma(K^0_L \to \pi^- l^+ \nu_l \gamma) + \Gamma(K^0_L \to \pi^+ l^- \bar{\nu}_l \gamma)$, which is measured experimentally (see also [4]). The results of calculation and experimental values for $K^0_L \to \pi^\pm l^\mp \nu_l \gamma$ branching ratios are presented in Table 1; here the following cuts in the kaon rest frame are used: $\omega \geqslant 30$ MeV and $\theta_{l\gamma} \geqslant 20^\circ$ (except the experimental results for $l = \mu$, where only restriction $\omega \geqslant 30$ MeV is imposed).

We use in calculation the particle masses $m_K = 497.6$ MeV, $m_\pi = 139.57$ MeV, $m_\mu = 105.658$ MeV, and $m_e = 0.511$ MeV, the Fermi coupling constant $G = 1.1664 \times 10^{-11}$ MeV^{-2}, the fine-structure constant $e^2/4\pi = \alpha = 1/137$, the Planck constant $\hbar = 6.582 \times 10^{-22}$ MeV·s, K^0_L mean life $\tau = 5.1 \times 10^{-8}$ s, $\sin\theta_c f_+(0) = 0.217$.

	$l = \mu$	$l = e$
Bijnens *et al.* [3]	5.2×10^{-4}	3.6×10^{-3}
present work	5.00×10^{-4}	3.45×10^{-3}
experimental values [9]	$(5.65 \pm 0.23) \times 10^{-4}$ $\omega \geqslant 30$ MeV	$(3.79 \pm 0.06) \times 10^{-3}$

Table 1: Branching ratios of $K^0_L \to \pi^\pm l^\mp \nu_l \gamma$ decays

The accuracy of our results can be estimated as follows. The leading corrections to them are due to the structure radiation from the hadronic vertex. They are proportional to the photon field strength, i.e. are on the order of ω. There are good reasons to believe that these corrections are less than the Low structure amplitudes which are on the order of ω^0. The Low contributions (including of course their interference with the accompanying radiation) to the discussed $K^0_L \to \pi^\pm \mu^\mp \nu_\mu \gamma$ and $K^0_L \to \pi^\pm e^\mp \nu_e \gamma$ branching ratios, according to our calculations, constitute 0.79×10^{-4} and 0.27×10^{-3}, respectively. Let us note also that corrections to the quoted results derived in Ref. [3] in the chiral perturbation theory are of similar magnitude.

As mentioned, additional corrections on the level of 20% to the branching ratios originate from our neglect of the form factor $f_-(t)$ and of the t-dependence of $f_+(t)$ (see also Table I in Ref. [2]). As to the relative accuracy of our numerical integration over phase space of final particles, it is about 10^{-3}.

To compare properly our results with those of Ref. [3] one should keep in mind that now the experimental values of some quantities are known with better accuracy. Indeed, we use $\sin\theta_c f_+(0) = 0.217$ in our calculation and, as far as we can see, in Ref. [3] the corresponding value is 0.22. Substitution of one of these values for another alters the results by about 3%.

Thus, our results for the branching ratios agree reasonably well with those of Ref. [3].

3. The T-odd triple momenta correlations $\xi = \vec{q}\cdot[\vec{p_l}\times\vec{p_\pi}]/M_K^3$ in the $K^0 \to \pi^- l^+ \nu_l \gamma$ decays arise from the interference term $2\mathrm{Re}(M_{tree}^\dagger A_{loop})$ in the decay rate; here M_{tree} is the tree amplitude and A_{loop} is the anti-Hermitian part of the loop diagrams presented below. As we are interested in the effect due to the electromagnetic final state interaction we consider only the one-loop diagrams generated from the tree ones by attaching the virtual photon. The on-mass-shell intermediate particles on the diagrams are marked by crosses.

The anti-Hermitian part of the sum of one-loop diagrams is written as

$$A_{loop} = \frac{i}{8\pi^2}\sum_n M_{fn}M_{in}^*, \tag{11}$$

where the sum over n includes the summation over the polarizations and the integration over the phase space of intermediate particles.

It is natural to divide the loop diagrams into four groups according to the type of amplitudes M_{fn} and M_{in}^*.

In the first group (see Fig. 2) the amplitude M_{fn} depicts the Compton scattering off the intermediate lepton (see Fig. 3):

$$M_{fn} = M_{3a} + M_{3b} = e^2\bar{v}_{k_l}\hat{e}_k\frac{\hat{P}-m_l}{2p_l q}\hat{e}^* v_l + e^2\bar{v}_{k_l}\hat{e}^*\frac{\hat{p}_l-\hat{k}-m_l}{-2p_l k}\hat{e}_k v_l. \tag{12}$$

As to M_{in}^*, it is the same tree amplitude (7), up to the change of some notations:

$$M_{in}^* = M_{ni} = M_{tree}\big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} = \frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\left\{(p_K+p_\pi)_\alpha\bar{u}_\nu\gamma^\alpha(1+\gamma_5)v_{k_l}\left(\frac{k_l e_k^*}{p_l q}-\frac{p_\pi e_k^*}{p_\pi k}\right)\right.$$
$$\left.+(p_K+p_\pi)_\alpha\bar{u}_\nu\gamma^\alpha(1+\gamma_5)\frac{\hat{k}\hat{e}_k^*}{2p_l q}v_{k_l} + \left(e_{k\alpha}^* - \frac{p_\pi e_k^*}{p_\pi k}k_\alpha\right)\bar{u}_\nu\gamma^\alpha(1+\gamma_5)v_{k_l}\right\}. \tag{13}$$

The element of phase space is

$$d\rho = \frac{d^3k}{2\omega_k}\frac{d^3k_l}{2\omega_l}\delta^{(4)}(k_l+k-P), \tag{14}$$



Figure 2: The one-loop diagrams (group I – the Compton scattering off the lepton)



a b

Figure 3: The Compton scattering off the lepton

where $P = p_l + q$.

In the second group of one-loop diagrams (see Fig. 4) the amplitude M_{fn} corresponds to the Compton scattering off the π-meson (see Fig. 5):

$$M_{fn} = M_{5a} + M_{5b} + M_{5c} = 2e^2 \left\{ -\frac{(p_\pi e^*)(k_\pi e_k)}{p_\pi q} + \frac{(p_\pi e_k)(k_\pi e^*)}{p_\pi k} + (e_k e^*) \right\}. \tag{15}$$

The amplitude M^*_{in} is the tree amplitude (7) up to the change of notations:

$$M^*_{in} = M_{ni} = M_{tree}\big|_{\substack{q\to k,\\ p_\pi \to k_\pi,\\ e\to e_k}} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0) \left\{ (p_K + k_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l \left(\frac{p_l e^*_k}{p_l k} - \frac{k_\pi e^*_k}{p_\pi q} \right) \right.$$

$$\left. + (p_K + k_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) \frac{\hat{k}\hat{e}^*_k}{2 p_l k} v_l + \left(e^*_{k\alpha} - \frac{k_\pi e^*_k}{p_\pi q} k_\alpha \right) \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_l \right\}. \tag{16}$$

The element of phase space is

$$d\rho' = \frac{d^3k}{2\omega_k} \frac{d^3k_\pi}{2\omega_\pi} \delta^{(4)}(k_\pi + k - P'), \tag{17}$$

where $P' = p_\pi + q$.



Figure 4: The one-loop diagrams (group II – the Compton scattering off the π-meson)

In the third group of one-loop diagrams (see Fig. 6) the amplitude M_{fn} is π-l scattering amplitude (see Fig. 7):

$$M_{fn} = M_7 = e^2 |F_\pi(k^2)| \frac{1}{k^2} (p_\pi + k_\pi)_\alpha \bar{v}_{k_l} \gamma^\alpha v_l. \tag{18}$$

The amplitude M_{in}^* is also the tree amplitude (7) up to the change of notations:

$$M_{in}^* = M_{ni} = M_{tree}|_{\substack{p_l \to k_l, \\ p_\pi \to k_\pi}} = \frac{G}{\sqrt{2}} \sin\theta_c e f_+(0) \left\{ (p_K + k_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_{k_l} \left(\frac{k_l e^*}{k_l q} - \frac{k_\pi e^*}{k_\pi q} \right) \right.$$
$$\left. + (p_K + k_\pi)_\alpha \bar{u}_\nu \gamma^\alpha (1+\gamma_5) \frac{\hat{q}\hat{e}^*}{2k_l q} v_{k_l} + \left(e_\alpha^* - \frac{k_\pi e^*}{k_\pi q} q_\alpha \right) \bar{u}_\nu \gamma^\alpha (1+\gamma_5) v_{k_l} \right\}. \tag{19}$$

The element of phase space is

$$d\rho'' = \frac{d^3 k_l}{2\omega_l} \frac{d^3 k_\pi}{2\omega_\pi} \delta^{(4)}(k_l + k_\pi - P''), \tag{20}$$

where $P'' = p_l + p_\pi$.

And finally, in the fourth group of one-loop diagrams (see Fig. 8) the amplitude M_{fn} is π-l scattering with emission of a photon (see Fig. 9):



a b c

Figure 5: The Compton scattering off the π-meson



Figure 6: The one-loop diagrams (group III – π-l scattering)

$$
\begin{aligned}
M_{fn} = M_{9a}+M_{9b}+M_{9c}+M_{9d}+M_{9e} = e^3|F_\pi(k^2)|\frac{1}{k^2}\Bigg\{&(p_\pi+k_\pi)_\alpha\bar{v}_{k_l}\gamma^\alpha\left(\frac{p_l e^*}{p_l q}+\frac{\hat{q}\hat{e}^*}{2p_l q}\right)v_l \\
-(p_\pi+k_\pi)_\alpha\bar{v}_{k_l}\left(\frac{k_l e^*}{k_l q}+\frac{\hat{q}\hat{e}^*}{2k_l q}\right)\gamma^\alpha v_l &- (p_\pi+k_\pi+q)_\alpha\bar{v}_{k_l}\gamma^\alpha v_l\frac{p_\pi e^*}{p_\pi q} \\
+(p_\pi+k_\pi-q)_\alpha\bar{v}_{k_l}\gamma^\alpha v_l\frac{k_\pi e^*}{k_\pi q} &+ 2e^*_\alpha\bar{v}_{k_l}\gamma^\alpha v_l\Bigg\}. \qquad (21)
\end{aligned}
$$

As to M^*_{in}, it is the amplitude of K^0_{l3} decays (see Fig. 10):

$$
M^*_{in} = M_{ni} = M_{10} = \frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)(p_K+k_\pi)_\alpha\bar{u}_\nu\gamma^\alpha(1+\gamma_5)v_{k_l}. \qquad (22)
$$

The element of phase space is

$$
d\rho''' = \frac{d^3k_l}{2\omega_l}\frac{d^3k_\pi}{2\omega_\pi}\delta^{(4)}(k_l+k_\pi-P'''), \qquad (23)
$$

where $P''' = p_l + p_\pi + q$.

In the discussed decays $k^2 < 0$ and $\sqrt{-k^2} < m_K(1-m_\pi^2/m_K^2) \lesssim 450$ MeV. At these energies the pion electromagnetic form factor can be approximated as

$$
|F_\pi(k^2)| \approx 1 + < r_\pi^2 > k^2/6, \qquad (24)
$$

here $< r_\pi^2 >$ is the pion mean square charge radius. According to our calculations (we use the PDG value $< r_\pi^2 >= (0.67\,\mathrm{fm})^2$ [9]), the contribution of term $< r_\pi^2 > k^2/6$ to the momenta correlation is small, namely, it does not exceed 5% of the contribution of 1. Therefore, we can put $|F_\pi(k^2)| = 1$. Moreover, keeping $< r_\pi^2 >$ term itself is beyond the accuracy of our calculation, because we have already neglected the form factor $f_-(t)$ and the t-dependence of $f_+(t)$.



Figure 7: The π-l scattering diagram



Figure 8: The one-loop diagrams (group IV – π-l scattering with emission of a photon)

The one-loop diagrams from group III contain infrared divergent terms, which can be omitted because infrared corrections to the decay width are factorized and, therefore, do not contribute to the effect.

The details of the calculation of the T-odd correlation $\xi = \vec{q} \cdot [\vec{p_l} \times \vec{p_\pi}]/M_K^3$ can be found in Appendices.

In fact, what is really measured experimentally is not the T-odd triple momenta correlation ξ by itself but the asymmetry

$$\begin{aligned} A_\xi = \frac{N_+ - N_-}{N_+ + N_-} &= \frac{\int (|M|^2_{even} + |M|^2_{odd})\, d\Phi_{\xi>0} - \int (|M|^2_{even} + |M|^2_{odd})\, d\Phi_{\xi<0}}{\int (|M|^2_{even} + |M|^2_{odd})\, d\Phi_{\xi>0} + \int (|M|^2_{even} + |M|^2_{odd})\, d\Phi_{\xi<0}} \\ &= \frac{\int |M|^2_{odd} d\Phi_{\xi>0}}{\int |M|^2_{even} d\Phi_{\xi>0}} \approx \frac{\int |M|^2_{odd} d\Phi_{\xi>0}}{\int |M_{tree}|^2 d\Phi_{\xi>0}} \end{aligned} \tag{25}$$

induced by this correlation; here N_+ and N_- are the numbers of events with $\xi > 0$ and $\xi < 0$, $|M|^2_{even}$ and $|M|^2_{odd}$ are the ξ-even and ξ-odd terms in the amplitude squared, and integration is performed over the phase space of the final particles.

The results for the asymmetry A_ξ in $K^0 \to \pi^- l^+ \nu_l \gamma$ decays are presented in Table 2 (here K^0 means both K_L^0 and K_S^0, and the following cuts in the kaon rest frame are used: $\omega \geqslant 30$ MeV and $\theta_{l\gamma} \geqslant 20°$).

The contributions of the Low term M_{Low} to the asymmetry A_ξ in $K^0 \to \pi^- \mu^+ \nu_\mu \gamma$ and $K^0 \to \pi^- e^+ \nu_e \gamma$ decays, according to our calculations, constitute 0.9×10^{-4} and -0.9×10^{-4}, respectively. Therefore, the contribution of M_{Low} for $K^0_{e3\gamma}$ decay is about 20%.



Figure 9: π-l scattering with emission of a photon



Figure 10: K^0_{l3} decays

For $K^0_{\mu 3\gamma}$ decay, however, it is large, comparable to the contributions of the accompanying radiation which are on the order of ω^{-1} and ω^0. So, it is difficult to estimate reliably the relative magnitude of the structure radiation contribution proportional to ω, i.e. to estimate reliably the true accuracy of thus-derived result for A_ξ in $K^0 \to \pi^-\mu^+\nu_\mu\gamma$ decay. However, we note here that corrections to the discussed asymmetry in the $K^+ \to \pi^0 l^+\nu_l\gamma$ decays, derived in Ref. [6] within the chiral perturbation theory, are small, on the order of 1%.

The asymmetry A_ξ was measured only in the $K^+_{l3\gamma}$ decays [11, 12]. The experimental errors are on the level of 10^{-1} for $l = \mu$ and 10^{-2} for $l = e$. Therefore, at present it is impossible to test the discussed theoretical predictions that are on the order of 10^{-4} [5, 7].

In any case experimental study of the T-odd asymmetry A_ξ in the $K_{l3\gamma}$ decays would be very important. It would allow one either to test CP-odd effects beyond the Standard Model or to set limits on them.

Acknowledgements

I am grateful to I.B. Khriplovich for his interest in the work and useful discussions as well as for critical reading of the text, to L.V. Kardapoltsev for his advice concerning numerical calculations, and to R.N. Lee and V.S. Fadin for pointing out the necessity to check the initial results.

	$l=\mu$	$l=e$
group I ($l-\gamma$)	-0.54×10^{-4}	-1.32×10^{-4}
group II ($\pi-\gamma$)	-3.6×10^{-4}	-3.2×10^{-4}
group III ($\pi-l$)	1.73×10^{-3}	8.6×10^{-4}
group IV ($\pi-l-\gamma$)	-1.41×10^{-3}	-8.6×10^{-4}
total	-1×10^{-4}	-4.5×10^{-4}

Table 2: A_ξ in $K^0\to\pi^-l^+\nu_l\gamma$ decays
($\omega\geqslant 30$ MeV and $\theta_{l\gamma}\geqslant 20^\circ$)

The work was supported in part by the Russian Foundation for Basic Research through Grant No. 11-02-00792-a, by the Federal Program "Personnel of Innovational Russia" through Grant No. 14.740.11.0082, and by Dmitry Zimin's Dynasty Foundation.

Appendix A

In this Appendix the list of the integrals that contribute to A_{loop} is given.
In formulas below we use the following notations:

$$I_{l\pi}=\sqrt{(p_lp_\pi)^2-m_l^2m_\pi^2},$$

$$I_{\pi P}=\sqrt{(p_\pi P)^2-m_\pi^2P^2},$$

$$I'_{lP}=\sqrt{(p_lP')^2-m_l^2P'^2},$$

$$I_{l\pi q}=\sqrt{(p_lp_\pi+p_lq+p_\pi q)^2-m_l^2m_\pi^2}.$$

We start with the integrals that contribute to group I of the one-loop diagrams:

$$\int d\rho=a_0=\frac{\pi(p_lq)}{P^2}; \tag{A.1}$$

$$\int k_\mu d\rho=a_PP_\mu,\ \text{where}\ \ a_P=\frac{p_lq}{P^2}a_0; \tag{A.2}$$

$$\int\frac{d\rho}{p_lk}=b_0=\frac{\pi}{2p_lq}\ln\left(\frac{P^2}{m_l^2}\right); \tag{A.3}$$

$$\int\frac{k_\mu}{p_lk}d\rho=B_{1\mu}=b_lp_{l\mu}+b_PP_\mu\,, \tag{A.4}$$

where b_l and b_P are the solutions of the set of equations

$$\begin{aligned}&b_lm_l^2+b_P(p_lP)=a_0,\\&b_l(p_lP)+b_PP^2=b_0(p_lq);\end{aligned}$$

$$\int \frac{k_\mu k_\nu}{p_l k} d\rho = B_{2\mu\nu} = b_2 g_{\mu\nu} + b_{ll} p_{l\mu} p_{l\nu} + b_{PP} P_\mu P_\nu + b_{lP}(p_{l\mu} P_\nu + P_\mu p_{l\nu}), \tag{A.5}$$

where b_2, b_{ll}, b_{PP}, and b_{lP} are the solutions of the set of equations

$$\begin{aligned} 4b_2 + b_{ll} m_l^2 + b_{PP} P^2 + 2b_{lP}(p_l P) &= 0, \\ b_2 + b_{ll} m_l^2 + b_{lP}(p_l P) &= 0, \\ b_{PP}(p_l P) + b_{lP} m_l^2 &= a_P, \\ b_2 + b_{PP} P^2 + b_{lP}(p_l P) &= b_P(p_l q); \end{aligned}$$

$$\int \frac{d\rho}{p_\pi k} = c_0 = \frac{\pi}{2 I_{\pi P}} \ln\left(\frac{p_\pi P + I_{\pi P}}{p_\pi P - I_{\pi P}} \right); \tag{A.6}$$

$$\int \frac{k_\mu}{p_\pi k} d\rho = C_{1\mu} = c_\pi p_{\pi\mu} + c_P P_\mu, \tag{A.7}$$

where c_π and c_P are the solutions of the set of equations

$$\begin{aligned} c_\pi m_\pi^2 + c_P(p_\pi P) &= a_0, \\ c_\pi(p_\pi P) + c_P P^2 &= c_0(p_l q); \end{aligned}$$

$$\int \frac{d\rho}{(p_l k)(p_\pi k)} = d_0 = \frac{\pi}{2(p_l q) I_{l\pi}} \ln\left(\frac{p_l p_\pi + I_{l\pi}}{p_l p_\pi - I_{l\pi}} \right); \tag{A.8}$$

$$\int \frac{k_\mu}{(p_l k)(p_\pi k)} d\rho = D_{1\mu} = d_l p_{l\mu} + d_\pi p_{\pi\mu} + d_P P_\mu, \tag{A.9}$$

where d_l, d_π, and d_P are the solutions of the set of equations

$$\begin{aligned} d_l m_l^2 + d_\pi(p_l p_\pi) + d_P(p_l P) &= c_0, \\ d_l(p_l p_\pi) + d_\pi m_\pi^2 + d_P(p_\pi P) &= b_0, \\ d_l(p_l P) + d_\pi(p_\pi P) + d_P P^2 &= d_0(p_l q); \end{aligned}$$

$$\begin{aligned} \int \frac{k_\mu k_\nu}{(p_l k)(p_\pi k)} d\rho = D_{2\mu\nu} &= d_2 g_{\mu\nu} + d_{ll} p_{l\mu} p_{l\nu} + d_{\pi\pi} p_{\pi\mu} p_{\pi\nu} + d_{PP} P_\mu P_\nu \\ &+ d_{l\pi}(p_{l\mu} p_{\pi\nu} + p_{\pi\mu} p_{l\nu}) + d_{lP}(p_{l\mu} P_\nu + P_\mu p_{l\nu}) + d_{\pi P}(p_{\pi\mu} P_\nu + P_\mu p_{\pi\nu}), \end{aligned} \tag{A.10}$$

where d_2, d_{ll}, $d_{\pi\pi}$, d_{PP}, $d_{l\pi}$, d_{lP}, and $d_{\pi P}$ are the solutions of the set of equations

$$\begin{aligned} 4d_2 + d_{ll} m_l^2 + d_{\pi\pi} m_\pi^2 + d_{PP} P^2 + 2d_{l\pi}(p_l p_\pi) + 2d_{lP}(p_l P) + 2d_{\pi P}(p_\pi P) &= 0, \\ d_2 + d_{ll} m_l^2 + d_{l\pi}(p_l p_\pi) + d_{lP}(p_l P) &= 0, \\ d_{\pi\pi}(p_l p_\pi) + d_{l\pi} m_l^2 + d_{\pi P}(p_l P) &= c_\pi, \\ d_{PP}(p_l P) + d_{lP} m_l^2 + d_{\pi P}(p_l p_\pi) &= c_P, \\ d_2 + d_{\pi\pi} m_\pi^2 + d_{l\pi}(p_l p_\pi) + d_{\pi P}(p_\pi P) &= 0, \end{aligned}$$

$$\begin{aligned} d_{PP}(p_\pi P) + d_{lP}(p_l p_\pi) + d_{\pi P} m_\pi^2 &= b_P, \\ d_2 + d_{PP}P^2 + d_{lP}(p_l P) + d_{\pi P}(p_\pi P) &= d_P(p_l q). \end{aligned}$$

Note that if in these integrals and sets of equations one changes all indices π (related to π-meson) to indices K (related to K-meson), then one gets integrals and sets of equations arising in calculation of T-odd triple momenta correlations in $K^+_{l3\gamma}$ decays [5, 7].

Most of the integrals that contribute to group II of the one-loop diagrams are those contributing to group I up to the change of some notations: indices l related to lepton must be changed by indices π related to π-meson and vice versa.

$$\int d\rho' = a'_0 = \frac{\pi(p_\pi q)}{P'^2}; \tag{A.11}$$

$$\int k_\mu d\rho' = a'_P P'_\mu, \text{ where } a'_P = \frac{p_\pi q}{P'^2} a'_0; \tag{A.12}$$

$$\int k_\mu k_\nu d\rho' = A'_{2\mu\nu} = a'_2 g_{\mu\nu} + a'_{PP} P'_\mu P'_\nu\,, \tag{A.13}$$

where a'_2 and a'_{PP} are the solutions of the set of equations

$$\begin{aligned} 4a'_2 + a'_{PP}P'^2 &= 0, \\ a'_2 + a'_{PP}P'^2 &= a'_P(p_\pi q); \end{aligned}$$

$$\int \frac{d\rho'}{p_\pi k} = b'_0 = \frac{\pi}{2p_\pi q} \ln\left(\frac{P'^2}{m_\pi^2}\right); \tag{A.14}$$

$$\int \frac{k_\mu}{p_\pi k} d\rho' = B'_{1\mu} = b'_\pi p_{\pi\mu} + b'_P P'_\mu\,, \tag{A.15}$$

where b'_π and b'_P are the solutions of the set of equations

$$\begin{aligned} b'_\pi m_\pi^2 + b'_P(p_\pi P') &= a'_0, \\ b'_\pi(p_\pi P') + b'_P P'^2 &= b'_0(p_\pi q); \end{aligned}$$

$$\int \frac{k_\mu k_\nu}{p_\pi k} d\rho' = B'_{2\mu\nu} = b'_2 g_{\mu\nu} + b'_{\pi\pi} p_{\pi\mu} p_{\pi\nu} + b'_{PP} P'_\mu P'_\nu + b'_{\pi P}(p_{\pi\mu} P'_\nu + P'_\mu p_{\pi\nu}), \tag{A.16}$$

where b', $b'_{\pi\pi}$, b'_{PP}, and $b'_{\pi P}$ are the solutions of the set of equations

$$\begin{aligned} 4b'_2 + b'_{\pi\pi} m_\pi^2 + b'_{PP}P'^2 + 2b'_{\pi P}(p_\pi P') &= 0, \\ b'_2 + b'_{\pi\pi} m_\pi^2 + b'_{\pi P}(p_\pi P') &= 0, \\ b'_{PP}(p_\pi P') + b'_{\pi P} m_\pi^2 &= a'_P, \\ b'_2 + b'_{PP}P'^2 + b'_{\pi P}(p_\pi P') &= b'_P(p_\pi q); \end{aligned}$$

$$\int \frac{d\rho'}{p_l k} = c'_0 = \frac{\pi}{2I'_{lP}} \ln\left(\frac{p_l P' + I'_{lP}}{p_l P' - I'_{lP}}\right); \tag{A.17}$$

$$\int \frac{k_\mu}{p_l k} d\rho' = C'_{1\mu} = c'_l p_{l\mu} + c'_P P'_\mu, \tag{A.18}$$

where c'_l and c'_P are the solutions of the set of equations

$$\begin{aligned} c'_l m_l^2 + c'_P (p_l P') &= a'_0, \\ c'_l (p_l P') + c'_P P'^2 &= c'_0 (p_\pi q); \end{aligned}$$

$$\int \frac{d\rho'}{(p_l k)(p_\pi k)} = d'_0 = \frac{\pi}{2(p_\pi q) I_{l\pi}} \ln\left(\frac{p_l p_\pi + I_{l\pi}}{p_l p_\pi - I_{l\pi}}\right); \tag{A.19}$$

$$\int \frac{k_\mu}{(p_l k)(p_\pi k)} d\rho' = D'_{1\mu} = d'_l p_{l\mu} + d'_\pi p_{\pi\mu} + d'_P P'_\mu, \tag{A.20}$$

where d'_l, d'_π, and d'_P are the solutions of the set of equations

$$\begin{aligned} d'_l m_l^2 + d'_\pi (p_l p_\pi) + d'_P (p_l P') &= b'_0, \\ d'_l (p_l p_\pi) + d'_\pi m_\pi^2 + d'_P (p_\pi P') &= c'_0, \\ d'_l (p_l P') + d'_\pi (p_\pi P') + d'_P P'^2 &= d'_0 (p_\pi q); \end{aligned}$$

$$\begin{aligned} \int \frac{k_\mu k_\nu}{(p_l k)(p_\pi k)} d\rho' = D'_{2\mu\nu} &= d'_2 g_{\mu\nu} + d'_{ll} p_{l\mu} p_{l\nu} + d'_{\pi\pi} p_{\pi\mu} p_{\pi\nu} + d'_{PP} P'_\mu P'_\nu \\ &+ d'_{l\pi}(p_{l\mu} p_{\pi\nu} + p_{\pi\mu} p_{l\nu}) + d'_{lP}(p_{l\mu} P'_\nu + P'_\mu p_{l\nu}) + d'_{\pi P}(p_{\pi\mu} P'_\nu + P'_\mu p_{\pi\nu}), \end{aligned} \tag{A.21}$$

where d'_2, d'_{ll}, $d'_{\pi\pi}$, d'_{PP}, $d'_{l\pi}$, d'_{lP}, and $d'_{\pi P}$ are the solutions of the set of equations

$$\begin{aligned} 4d'_2 + d'_{ll} m_l^2 + d'_{\pi\pi} m_\pi^2 + d'_{PP} P'^2 + 2d'_{l\pi}(p_l p_\pi) + 2d'_{lP}(p_l P') + 2d'_{\pi P}(p_\pi P') &= 0, \\ d'_2 + d'_{ll} m_l^2 + d'_{l\pi}(p_l p_\pi) + d'_{lP}(p_l P') &= 0, \\ d'_{\pi\pi}(p_l p_\pi) + d'_{l\pi} m_l^2 + d'_{\pi P}(p_l P') &= b'_\pi, \\ d'_{PP}(p_l P') + d'_{lP} m_l^2 + d'_{\pi P}(p_l p_\pi) &= b'_P, \\ d'_2 + d'_{\pi\pi} m_\pi^2 + d'_{l\pi}(p_l p_\pi) + d'_{\pi P}(p_\pi P') &= 0, \\ d'_{PP}(p_\pi P') + d'_{lP}(p_l p_\pi) + d'_{\pi P} m_\pi^2 &= c'_P, \\ d'_2 + d'_{PP} P'^2 + d'_{lP}(p_l P') + d'_{\pi P}(p_\pi P') &= d'_P (p_\pi q). \end{aligned}$$

The integrals arising in calculation of group III of the one-loop diagrams:

$$\int d\rho'' = a''_0 = \frac{\pi I_{l\pi}}{P''^2}; \tag{A.22}$$

$$\int k_{l\mu} d\rho'' = a''_P P''_\mu, \text{ where } a''_P = \frac{(p_l P'')}{P''^2} a''_0; \tag{A.23}$$

$$\int \frac{d\rho''}{k_l q} = b''_0 = \frac{\pi}{2P'' q} \ln\left(\frac{p_l P'' + I_{l\pi}}{p_l P'' - I_{l\pi}}\right); \tag{A.24}$$

$$\int \frac{k_{l\mu}}{k_l q} d\rho'' = B''_{1\mu} = b''_q q_\mu + b''_P P''_\mu, \tag{A.25}$$

where b''_q and b''_P are the solutions of the set of equations

$$\begin{aligned} b''_P(P''q) &= a''_0, \\ b''_q(P''q) + b''_P P''^2 &= b''_0(p_l P''); \end{aligned}$$

$$\int \frac{k_{l\mu}k_{l\nu}}{k_l q} d\rho'' = B''_{2\mu\nu} = b''_2 g_{\mu\nu} + b''_{qq} q_\mu q_\nu + b''_{PP} P''_\mu P''_\nu + b''_{qP}(q_\mu P''_\nu + P''_\mu q_\nu), \tag{A.26}$$

where b''_2, b''_{qq}, b''_{PP}, and b''_{qP} are the solutions of the set of equations

$$\begin{aligned} 4b''_2 + b''_{PP} P''^2 + 2b''_{qP}(P''q) &= m_l^2 b''_0, \\ b''_2 + b''_{qP}(P''q) &= 0, \\ b''_{qq}(P''q) + b''_{qP} P''^2 &= b''_q(p_l P''), \\ b''_2 + b''_{PP} P''^2 + b''_{qP}(P''q) &= b''_P(p_l P''); \end{aligned}$$

$$\int \frac{d\rho''}{k_\pi q} = c''_0 = \frac{\pi}{2P''q} \ln\left(\frac{p_\pi P'' + I_{l\pi}}{p_\pi P'' - I_{l\pi}}\right); \tag{A.27}$$

$$\int \frac{k_{\pi\mu}}{k_\pi q} d\rho'' = C''_{1\mu} = c''_q q_\mu + c''_P P''_\mu, \tag{A.28}$$

where c''_q and c''_P are the solutions of the set of equations

$$\begin{aligned} c''_P(P''q) &= a''_0, \\ c''_q(P''q) + c''_P P''^2 &= c''_0(p_\pi P''); \end{aligned}$$

$$\int \frac{k_{\pi\mu}k_{\pi\nu}}{k_\pi q} d\rho'' = C''_{2\mu\nu} = c''_2 g_{\mu\nu} + c''_{qq} q_\mu q_\nu + c''_{PP} P''_\mu P''_\nu + c''_{qP}(q_\mu P''_\nu + P''_\mu q_\nu), \tag{A.29}$$

where c''_2, c''_{qq}, c''_{PP}, and c''_{qP} are the solutions of the set of equations

$$\begin{aligned} 4c''_2 + c''_{PP} P''^2 + 2c''_{qP}(P''q) &= m_\pi^2 c''_0, \\ c''_2 + c''_{qP}(P''q) &= 0, \\ c''_{qq}(P''q) + c''_{qP} P''^2 &= c''_q(p_\pi P''), \\ c''_2 + c''_{PP} P''^2 + c''_{qP}(P''q) &= c''_P(p_\pi P''); \end{aligned}$$

$$\int \frac{d\rho''}{k^2} = \frac{\pi}{4I_{l\pi}} \ln\left(\frac{\mu^2 P''^2}{4I_{l\pi}^2}\right) \to 0, \tag{A.30}$$

here μ^2 is the cut-off parameter for the logarithmically divergent integral over k^2, and the sign $\to$ means that infrared divergent terms are omitted;

$$\int \frac{k_{l\mu}}{k^2} d\rho'' \to D''_{1\mu} = \frac{\pi}{2I_{l\pi}} \left(\frac{p_\pi P''}{P''^2} p_{l\mu} - \frac{p_l P''}{P''^2} p_{\pi\mu} \right); \tag{A.31}$$

$$\int \frac{d\rho''}{k^2(k_l q)} = \frac{\pi}{4(p_l q) I_{l\pi}} \left[\ln\left(\frac{\mu^2 P''^2}{4 I_{l\pi}^2}\right) + \ln\left(\frac{m_l^2 (P''q)^2}{P''^2 (p_l q)^2}\right)\right] \to e''_0 = \frac{\pi}{4(p_l q) I_{l\pi}} \ln\left(\frac{m_l^2 (P''q)^2}{P''^2 (p_l q)^2}\right); \tag{A.32}$$

$$\int \frac{k_{l\mu}}{k^2(k_l q)} d\rho'' \to E''_{1\mu} = e''_l p_{l\mu} + e''_\pi p_{\pi\mu} + e''_q q_\mu, \tag{A.33}$$

where e''_l, e''_π, and e''_q are the solutions of the set of equations

$$\begin{aligned} e''_l (p_l p_\pi) + e''_\pi m_\pi^2 + e''_q (p_\pi q) &= \frac{1}{2}\, b''_0 + (p_l p_\pi) e''_0, \\ e''_l m_l^2 + e''_\pi (p_l p_\pi) + e''_q (p_l q) &= -\frac{1}{2}\, b''_0 + m_l^2 e''_0, \\ e''_l (p_l q) + e''_\pi (p_\pi q) &= 0; \end{aligned}$$

$$\begin{aligned} \int \frac{k_{l\mu} k_{l\nu}}{k^2(k_l q)} d\rho'' \to E''_{2\mu\nu} &= e''_2 g_{\mu\nu} + e''_{ll} p_{l\mu} p_{l\nu} + e''_{\pi\pi} p_{\pi\mu} p_{\pi\nu} + e''_{qq} q_\mu q_\nu \\ &+ e''_{l\pi} (p_{l\mu} p_{\pi\nu} + p_{\pi\mu} p_{l\nu}) + e''_{lq} (p_{l\mu} q_\nu + q_\mu p_{l\nu}) + e''_{\pi q} (p_{\pi\mu} q_\nu + q_\mu p_{\pi\nu}), \end{aligned} \tag{A.34}$$

where e''_2, e''_{ll}, $e''_{\pi\pi}$, e''_{qq}, $e''_{l\pi}$, e''_{lq}, and $e''_{\pi q}$ are the solutions of the set of equations

$$\begin{aligned} 4e''_2 + e''_{ll} m_l^2 + e''_{\pi\pi} m_\pi^2 + 2e''_{l\pi} (p_l p_\pi) + 2e''_{lq} (p_l q) + 2e''_{\pi q} (p_\pi q) &= m_l^2 e''_0, \\ e''_2 m_l^2 + e''_{ll} m_l^4 + e''_{\pi\pi} (p_l p_\pi)^2 + e''_{qq} (p_l q)^2 + 2e''_{l\pi} m_l^2 (p_l p_\pi) + 2e''_{lq} m_l^2 (p_l q) + 2e''_{\pi q} (p_l p_\pi)(p_l q) & \\ = \frac{I_{l\pi}^2 (D''_1 q)}{(P''q)^2} - \frac{1}{2} \left(\frac{(p_l q)(p_l P'')}{P''q} + m_l^2 \right) b''_0 + m_l^4 e''_0, & \\ e''_2 m_\pi^2 + e''_{ll} (p_l p_\pi)^2 + e''_{\pi\pi} m_\pi^4 + e''_{qq} (p_\pi q)^2 + 2e''_{l\pi} m_\pi^2 (p_l p_\pi) + 2e''_{lq} (p_l p_\pi)(p_\pi q) + 2e''_{\pi q} m_\pi^2 (p_\pi q) & \\ = \frac{I_{l\pi}^2 (D''_1 q)}{(P''q)^2} + \frac{1}{2} \left(\frac{(p_\pi q)(p_l P'')}{P''q} + (p_l p_\pi) \right) b''_0 + (p_l p_\pi)^2 e''_0, & \\ e''_2 (p_l p_\pi) + e''_{ll} m_l^2 (p_l p_\pi) + e''_{\pi\pi} m_\pi^2 (p_l p_\pi) + e''_{qq} (p_l q)(p_\pi q) + e''_{l\pi} (m_l^2 m_\pi^2 + (p_l p_\pi)^2) & \\ + e''_{lq} ((p_l p_\pi)(p_l q) + m_l^2 (p_\pi q)) + e''_{\pi q} ((p_l p_\pi)(p_\pi q) + m_\pi^2 (p_l q)) & \\ = -\frac{I_{l\pi}^2 (D''_1 q)}{(P''q)^2} + \frac{1}{2} \left(m_l^2 - \frac{(p_\pi q)(p_l P'')}{P''q} \right) b''_0 + m_l^2 (p_l p_\pi) e''_0, & \\ e''_2 + e''_{lq} (p_l q) + e''_{\pi q} (p_\pi q) &= 0, \\ e''_{ll} (p_l q) + e''_{l\pi} (p_\pi q) &= \frac{\pi}{2I_{l\pi}} \frac{p_\pi P''}{P''^2}, \\ e''_{\pi\pi} (p_\pi q) + e''_{l\pi} (p_l q) &= -\frac{\pi}{2I_{l\pi}} \frac{p_l P''}{P''^2}; \end{aligned}$$

$$\int \frac{d\rho''}{k^2(k_\pi q)} = \frac{\pi}{4(p_\pi q)I_{l\pi}}\left[\ln\left(\frac{\mu^2 P''^2}{4I^2_{l\pi}}\right)+\ln\left(\frac{m_\pi^2(P''q)^2}{P''^2(p_\pi q)^2}\right)\right] \to f''_0 = \frac{\pi}{4(p_\pi q)I_{l\pi}}\ln\left(\frac{m_\pi^2(P''q)^2}{P''^2(p_\pi q)^2}\right); \tag{A.35}$$

$$\int \frac{k_{\pi\mu}}{k^2(k_\pi q)}d\rho'' \to F''_{1\mu} = f''_l p_{l\mu} + f''_\pi p_{\pi\mu} + f''_q q_\mu, \tag{A.36}$$

where f''_l, f''_π, and f''_q are the solutions of the set of equations

$$\begin{aligned}
f''_l m_l^2 + f''_\pi(p_l p_\pi) + f''_q(p_l q) &= \frac{1}{2}\, c''_0 + (p_l p_\pi) f''_0,\\
f''_l(p_l p_\pi) + f''_\pi m_\pi^2 + f''_q(p_\pi q) &= -\frac{1}{2}\, c''_0 + m_\pi^2 f''_0,\\
f''_l(p_l q) + f''_\pi(p_\pi q) &= 0;
\end{aligned}$$

$$\begin{aligned}
\int \frac{k_{\pi\mu}k_{\pi\nu}}{k^2(k_\pi q)}d\rho'' \to F''_{2\mu\nu} = {}& f''_2 g_{\mu\nu} + f''_{ll}p_{l\mu}p_{l\nu} + f''_{\pi\pi}p_{\pi\mu}p_{\pi\nu} + f''_{qq}q_\mu q_\nu\\
& + f''_{l\pi}(p_{l\mu}p_{\pi\nu} + p_{\pi\mu}p_{l\nu}) + f''_{lq}(p_{l\mu}q_\nu + q_\mu p_{l\nu}) + f''_{\pi q}(p_{\pi\mu}q_\nu + q_\mu p_{\pi\nu}),
\end{aligned} \tag{A.37}$$

where f''_2, f''_{ll}, $f''_{\pi\pi}$, f''_{qq}, $f''_{l\pi}$, f''_{lq}, and $f''_{\pi q}$ are the solutions of the set of equations

$$\begin{aligned}
4f''_2 + f''_{ll}m_l^2 + f''_{\pi\pi}m_\pi^2 + 2f''_{l\pi}(p_l p_\pi) + 2f''_{lq}(p_l q) + 2f''_{\pi q}(p_\pi q) &= m_\pi^2 f''_0,\\
f''_2 m_l^2 + f''_{ll}m_l^4 + f''_{\pi\pi}(p_l p_\pi)^2 + f''_{qq}(p_l q)^2 + 2f''_{l\pi}m_l^2(p_l p_\pi) + 2f''_{lq}m_l^2(p_l q) + 2f''_{\pi q}(p_l p_\pi)(p_l q)&\\
= -\frac{I^2_{l\pi}(D''_1 q)}{(P''q)^2} + \frac{1}{2}\left(\frac{(p_l q)(p_\pi P'')}{P''q} + p_l p_\pi\right) c''_0 + (p_l p_\pi)^2 f''_0&,\\
f''_2 m_\pi^2 + f''_{ll}(p_l p_\pi)^2 + f''_{\pi\pi}m_\pi^4 + f''_{qq}(p_\pi q)^2 + 2f''_{l\pi}m_\pi^2(p_l p_\pi) + 2f''_{lq}(p_l p_\pi)(p_\pi q) + 2f''_{\pi q}m_\pi^2(p_\pi q)&\\
= -\frac{I^2_{l\pi}(D''_1 q)}{(P''q)^2} - \frac{1}{2}\left(\frac{(p_\pi q)(p_\pi P'')}{P''q} + m_\pi^2\right) c''_0 + m_\pi^4 f''_0&,\\
f''_2(p_l p_\pi) + f''_{ll}m_l^2(p_l p_\pi) + f''_{\pi\pi}m_\pi^2(p_l p_\pi) + f''_{qq}(p_l q)(p_\pi q) + f''_{l\pi}(m_l^2 m_\pi^2 + (p_l p_\pi)^2)&\\
+f''_{lq}((p_l p_\pi)(p_l q) + m_l^2(p_\pi q)) + f''_{\pi q}((p_l p_\pi)(p_\pi q) + m_\pi^2(p_l q))&\\
= \frac{I^2_{l\pi}(D''_1 q)}{(P''q)^2} + \frac{1}{2}\left(m_\pi^2 - \frac{(p_l q)(p_\pi P'')}{P''q}\right) c''_0 + m_\pi^2(p_l p_\pi) f''_0&,\\
f''_2 + f''_{lq}(p_l q) + f''_{\pi q}(p_\pi q) &= 0,\\
f''_{ll}(p_l q) + f''_{l\pi}(p_\pi q) &= -\frac{\pi}{2I_{l\pi}}\frac{p_\pi P''}{P''^2},\\
f''_{\pi\pi}(p_\pi q) + f''_{l\pi}(p_l q) &= \frac{\pi}{2I_{l\pi}}\frac{p_l P''}{P''^2}.
\end{aligned}$$

The integrals arising in calculation of group IV of the one-loop diagrams:

$$\int d\rho''' = a'''_0 = \frac{\pi I_{l\pi q}}{P'''^2}; \tag{A.38}$$

$$\int k_{l\mu} d\rho''' = a'''_P P'''_\mu, \text{ where } a'''_P = \frac{p_l P''' + p_\pi q}{P'''^2} a'''_0; \tag{A.39}$$

$$\int \frac{d\rho'''}{k_l q} = b'''_0 = \frac{\pi}{2P''q} \ln\left(\frac{p_l P''' + p_\pi q + I_{l\pi q}}{p_l P''' + p_\pi q - I_{l\pi q}} \right); \tag{A.40}$$

$$\int \frac{k_{l\mu}}{k_l q} d\rho''' = B'''_{1\mu} = b'''_q q_\mu + b'''_P P'''_\mu, \tag{A.41}$$

where b'''_q and b'''_P are the solutions of the set of equations

$$\begin{aligned} b'''_P(P''q) &= a'''_0, \\ b'''_q(P''q) + b'''_P P'''^2 &= b'''_0(p_l P''' + p_\pi q); \end{aligned}$$

$$\int \frac{k_{l\mu} k_{l\nu}}{k_l q} d\rho''' = B'''_{2\mu\nu} = b'''_2 g_{\mu\nu} + b'''_{qq} q_\mu q_\nu + b'''_{PP} P'''_\mu P'''_\nu + b'''_{qP}(q_\mu P'''_\nu + P'''_\mu q_\nu), \tag{A.42}$$

where b'''_2, b'''_{qq}, b'''_{PP}, and b'''_{qP} are the solutions of the set of equations

$$\begin{aligned} 4b'''_2 + b'''_{PP} P'''^2 + 2b'''_{qP}(P''q) &= m_l^2 b'''_0, \\ b'''_2 + b'''_{qP}(P''q) &= 0, \\ b'''_{qq}(P''q) + b'''_{qP} P'''^2 &= b'''_q(p_l P''' + p_\pi q), \\ b'''_2 + b'''_{PP} P'''^2 + b'''_{qP}(P''q) &= b'''_P(p_l P''' + p_\pi q); \end{aligned}$$

$$\int \frac{d\rho'''}{k_\pi q} = c'''_0 = \frac{\pi}{2P''q} \ln\left(\frac{p_\pi P''' + p_l q + I_{l\pi q}}{p_\pi P''' + p_l q - I_{l\pi q}} \right); \tag{A.43}$$

$$\int \frac{k_{\pi\mu}}{k_\pi q} d\rho''' = C'''_{1\mu} = c'''_q q_\mu + c'''_P P'''_\mu, \tag{A.44}$$

where c'''_q and c'''_P are the solutions of the set of equations

$$\begin{aligned} c'''_P(P''q) &= a'''_0, \\ c'''_q(P''q) + c'''_P P'''^2 &= c'''_0(p_\pi P''' + p_l q); \end{aligned}$$

$$\int \frac{k_{\pi\mu} k_{\pi\nu}}{k_\pi q} d\rho''' = C'''_{2\mu\nu} = c'''_2 g_{\mu\nu} + c'''_{qq} q_\mu q_\nu + c'''_{PP} P'''_\mu P'''_\nu + c'''_{qP}(q_\mu P'''_\nu + P'''_\mu q_\nu), \tag{A.45}$$

where c'''_2, c'''_{qq}, c'''_{PP}, and c'''_{qP} are the solutions of the set of equations

$$\begin{aligned} 4c'''_2 + c'''_{PP} P'''^2 + 2c'''_{qP}(P''q) &= m_\pi^2 c'''_0, \\ c'''_2 + c'''_{qP}(P''q) &= 0, \\ c'''_{qq}(P''q) + c'''_{qP} P'''^2 &= c'''_q(p_\pi P''' + p_l q), \\ c'''_2 + c'''_{PP} P'''^2 + c'''_{qP}(P''q) &= c'''_P(p_\pi P''' + p_l q); \end{aligned}$$

$$\int \frac{d\rho'''}{(p_\pi - k_\pi)^2} = d_0''' = \frac{\pi}{4I_{\pi P}} \ln\left(\frac{(p_\pi P)(p_l p_\pi + p_l q + p_\pi q) - m_\pi^2 (p_l P) - I_{l\pi q} I_{\pi P}}{(p_\pi P)(p_l p_\pi + p_l q + p_\pi q) - m_\pi^2 (p_l P) + I_{l\pi q} I_{\pi P}} \right); \tag{A.46}$$

$$\int \frac{k_{l\mu}}{(p_\pi - k_\pi)^2} d\rho''' = D_{1\mu}''' = d_\pi''' p_{\pi\mu} + d_P''' P_\mu, \tag{A.47}$$

where d_π''' and d_P''' are the solutions of the set of equations

$$\begin{aligned} d_\pi''' m_\pi^2 + d_P''' (p_\pi P) &= \frac{1}{2} a_0''' + (p_\pi P) d_0''', \\ d_\pi''' (p_\pi P) + d_P''' P^2 &= -\frac{1}{2} a_0''' + (p_l P) d_0'''; \end{aligned}$$

$$\int \frac{d\rho'''}{(p_l - k_l)^2} = e_0''' = \frac{\pi}{4I_{lP}'} \ln\left(\frac{(p_l P')(p_l p_\pi + p_l q + p_\pi q) - m_l^2 (p_\pi P') - I_{l\pi q} I_{lP}'}{(p_l P')(p_l p_\pi + p_l q + p_\pi q) - m_l^2 (p_\pi P') + I_{l\pi q} I_{lP}'} \right); \tag{A.48}$$

$$\int \frac{k_{\pi\mu}}{(p_l - k_l)^2} d\rho''' = E_{1\mu}''' = e_l''' p_{l\mu} + e_P''' P_\mu', \tag{A.49}$$

where e_l''' and e_P''' are the solutions of the set of equations

$$\begin{aligned} e_l''' m_l^2 + e_P''' (p_l P') &= \frac{1}{2} a_0''' + (p_l P') e_0''', \\ e_l''' (p_l P') + e_P''' P'^2 &= -\frac{1}{2} a_0''' + (p_\pi P') e_0'''; \end{aligned}$$

$$\int \frac{d\rho'''}{(p_\pi - k_\pi)^2 (k_l q)} = f_0''' = \frac{\pi}{4(p_l q) I_{l\pi}} \ln\left(\frac{p_l p_\pi (p_l p_\pi + p_l q + p_\pi q) - m_l^2 m_\pi^2 - I_{l\pi} I_{l\pi q}}{p_l p_\pi (p_l p_\pi + p_l q + p_\pi q) - m_l^2 m_\pi^2 + I_{l\pi} I_{l\pi q}} \right); \tag{A.50}$$

$$\int \frac{k_{l\mu}}{(p_\pi - k_\pi)^2 (k_l q)} d\rho''' = F_{1\mu}''' = f_l''' p_{l\mu} + f_\pi''' p_{\pi\mu} + f_q''' q_\mu, \tag{A.51}$$

where f_l''', f_π''', and f_q''' are the solutions of the set of equations

$$\begin{aligned} f_l''' m_l^2 + f_\pi''' (p_l p_\pi) + f_q''' (p_l q) &= -\frac{1}{2} b_0''' - d_0''' + (p_l P) f_0''', \\ f_l''' (p_l p_\pi) + f_\pi''' m_\pi^2 + f_q''' (p_\pi q) &= \frac{1}{2} b_0''' + (p_\pi P) f_0''', \\ f_l''' (p_l q) + f_\pi''' (p_\pi q) &= d_0'''; \end{aligned}$$

$$\begin{aligned} \int \frac{k_{l\mu} k_{l\nu}}{(p_\pi - k_\pi)^2 (k_l q)} d\rho''' = F_{2\mu\nu}''' &= f_2''' g_{\mu\nu} + f_{ll}''' p_{l\mu} p_{l\nu} + f_{\pi\pi}''' p_{\pi\mu} p_{\pi\nu} + f_{qq}''' q_\mu q_\nu \\ &+ f_{l\pi}''' (p_{l\mu} p_{\pi\nu} + p_{\pi\mu} p_{l\nu}) + f_{lq}''' (p_{l\mu} q_\nu + q_\mu p_{l\nu}) + f_{\pi q}''' (p_{\pi\mu} q_\nu + q_\mu p_{\pi\nu}), \end{aligned} \tag{A.52}$$

where f_2''', f_{ll}''', $f_{\pi\pi}'''$, f_{qq}''', $f_{l\pi}'''$, f_{lq}''', and $f_{\pi q}'''$ are the solutions of the set of equations

$$\begin{aligned}
4f_2''' + f_{ll}'''m_l^2 + f_{\pi\pi}'''m_\pi^2 + 2f_{l\pi}'''(p_lp_\pi) + 2f_{lq}'''(p_lq) + 2f_{\pi q}'''(p_\pi q) = m_l^2 f_0''', \\
f_2'''m_l^2 + f_{ll}'''m_l^4 + f_{\pi\pi}'''(p_lp_\pi)^2 + f_{qq}'''(p_lq)^2 + 2f_{l\pi}'''m_l^2(p_lp_\pi) + 2f_{lq}'''m_l^2(p_lq) + 2f_{\pi q}'''(p_lp_\pi)(p_lq) \\
= (p_lP)^2 f_0''' - \frac{1}{2}\left(p_lP + \frac{p_lq}{P''q}(p_lP''' + p_\pi q)\right) b_0''' - (p_lP)d_0''' \\
+\frac{\pi}{2P''q}\left(\frac{p_\pi P'''}{P'''^2} - \frac{p_\pi q}{P''q}\right) I_{l\pi q} + \frac{1}{2}a_0''' - (D_1'''p_l), \\
f_2'''m_\pi^2 + f_{ll}'''(p_lp_\pi)^2 + f_{\pi\pi}'''m_\pi^4 + f_{qq}'''(p_\pi q)^2 + 2f_{l\pi}'''m_\pi^2(p_lp_\pi) + 2f_{lq}'''(p_lp_\pi)(p_\pi q) + 2f_{\pi q}'''m_\pi^2(p_\pi q) \\
= (p_\pi P)^2 f_0''' + \frac{1}{2}\left(p_\pi P + \frac{p_\pi q}{P''q}(p_lP''' + p_\pi q)\right) b_0''' + \frac{\pi}{2P''q}\left(\frac{p_\pi P'''}{P'''^2} - \frac{p_\pi q}{P''q}\right) I_{l\pi q}, \\
f_2'''(p_lp_\pi) + f_{ll}'''m_l^2(p_lp_\pi) + f_{\pi\pi}'''m_\pi^2(p_lp_\pi) + f_{qq}'''(p_lq)(p_\pi q) + f_{l\pi}'''(m_l^2m_\pi^2 + (p_lp_\pi)^2) \\
+f_{lq}'''((p_lp_\pi)(p_lq) + m_l^2(p_\pi q)) + f_{\pi q}'''((p_lp_\pi)(p_\pi q) + m_\pi^2(p_lq)) \\
= (p_lP)(p_\pi P)f_0''' + \frac{1}{2}\left(p_lP - \frac{p_\pi q}{P''q}(p_lP''' + p_\pi q)\right) b_0''' - (p_\pi P)d_0''' \\
-\frac{1}{2}a_0''' - \frac{\pi}{2P''q}\left(\frac{p_\pi P'''}{P'''^2} - \frac{p_\pi q}{P''q}\right) I_{l\pi q}, \\
f_2''' + f_{lq}'''(p_lq) + f_{\pi q}'''(p_\pi q) = d_P''', \\
f_{ll}'''(p_lq) + f_{l\pi}'''(p_\pi q) = d_P''', \\
f_{\pi\pi}'''(p_\pi q) + f_{l\pi}'''(p_lq) = d_\pi''';
\end{aligned}$$

$$\int \frac{d\rho'''}{(p_l - k_l)^2(k_\pi q)} = g_0''' = \frac{\pi}{4(p_\pi q)I_{l\pi}} \ln\left(\frac{p_lp_\pi(p_lp_\pi + p_lq + p_\pi q) - m_l^2m_\pi^2 - I_{l\pi}I_{l\pi q}}{p_lp_\pi(p_lp_\pi + p_lq + p_\pi q) - m_l^2m_\pi^2 + I_{l\pi}I_{l\pi q}}\right); \tag{A.53}$$

$$\int \frac{k_{\pi\mu}}{(p_l - k_l)^2(k_\pi q)} d\rho''' = G_{1\mu}''' = g_l'''p_{l\mu} + g_\pi'''p_{\pi\mu} + g_q'''q_\mu, \tag{A.54}$$

where g_l''', g_π''', and g_q''' are the solutions of the set of equations

$$\begin{aligned}
g_l'''m_l^2 + g_\pi'''(p_lp_\pi) + g_q'''(p_lq) &= \frac{1}{2}c_0''' + (p_lP')g_0''', \\
g_l'''(p_lp_\pi) + g_\pi'''m_\pi^2 + g_q'''(p_\pi q) &= -\frac{1}{2}c_0''' - e_0''' + (p_\pi P')g_0''', \\
g_l'''(p_lq) + g_\pi'''(p_\pi q) &= e_0''';
\end{aligned}$$

$$\begin{aligned}
\int \frac{k_{\pi\mu}k_{\pi\nu}}{(p_l - k_l)^2(k_\pi q)} d\rho''' = G_{2\mu\nu}''' = g_2'''g_{\mu\nu} + g_{ll}'''p_{l\mu}p_{l\nu} + g_{\pi\pi}'''p_{\pi\mu}p_{\pi\nu} + g_{qq}'''q_\mu q_\nu \\
+ g_{l\pi}'''(p_{l\mu}p_{\pi\nu} + p_{\pi\mu}p_{l\nu}) + g_{lq}'''(p_{l\mu}q_\nu + q_\mu p_{l\nu}) + g_{\pi q}'''(p_{\pi\mu}q_\nu + q_\mu p_{\pi\nu}),
\end{aligned} \tag{A.55}$$

where g_2''', g_{ll}''', $g_{\pi\pi}'''$, g_{qq}''', $g_{l\pi}'''$, g_{lq}''', and $g_{\pi q}'''$ are the solutions of the set of equations

$$\begin{aligned}
4g_2''' + g_{ll}'''m_l^2 + g_{\pi\pi}'''m_\pi^2 + 2g_{l\pi}'''(p_lp_\pi) + 2g_{lq}'''(p_lq) + 2g_{\pi q}'''(p_\pi q) = m_\pi^2 g_0''', \\
g_2'''m_l^2 + g_{ll}'''m_l^4 + g_{\pi\pi}'''(p_lp_\pi)^2 + g_{qq}'''(p_lq)^2 + 2g_{l\pi}'''m_l^2(p_lp_\pi) + 2g_{lq}'''m_l^2(p_lq) + 2g_{\pi q}'''(p_lp_\pi)(p_lq) \\
= (p_lP')^2 g_0''' + \frac{1}{2}\left(p_lP' + \frac{p_lq}{P''q}(p_\pi P''' + p_lq)\right) c_0''' + \frac{\pi}{2P''q}\left(\frac{p_lP'''}{P'''^2} - \frac{p_lq}{P''q}\right) I_{l\pi q}, \\
g_2'''m_\pi^2 + g_{ll}'''(p_lp_\pi)^2 + g_{\pi\pi}'''m_\pi^4 + g_{qq}'''(p_\pi q)^2 + 2g_{l\pi}'''m_\pi^2(p_lp_\pi) + 2g_{lq}'''(p_lp_\pi)(p_\pi q) + 2g_{\pi q}'''m_\pi^2(p_\pi q) \\
= (p_\pi P')^2 g_0''' - \frac{1}{2}\left(p_\pi P' + \frac{p_\pi q}{P''q}(p_\pi P''' + p_lq)\right) c_0''' - (p_\pi P')e_0''' \\
+\frac{\pi}{2P''q}\left(\frac{p_lP'''}{P'''^2} - \frac{p_lq}{P''q}\right) I_{l\pi q} + \frac{1}{2}a_0''' - (E_1'''p_\pi), \\
g_2'''(p_lp_\pi) + g_{ll}'''m_l^2(p_lp_\pi) + g_{\pi\pi}'''m_\pi^2(p_lp_\pi) + g_{qq}'''(p_lq)(p_\pi q) + g_{l\pi}'''(m_l^2m_\pi^2 + (p_lp_\pi)^2) \\
+g_{lq}'''((p_lp_\pi)(p_lq) + m_l^2(p_\pi q)) + g_{\pi q}'''((p_lp_\pi)(p_\pi q) + m_\pi^2(p_lq)) \\
= (p_\pi P')(p_lP')g_0''' + \frac{1}{2}\left(p_\pi P' - \frac{p_lq}{P''q}(p_\pi P''' + p_lq)\right) c_0''' - (p_lP')e_0''' \\
-\frac{1}{2}a_0''' - \frac{\pi}{2P''q}\left(\frac{p_lP'''}{P'''^2} - \frac{p_lq}{P''q}\right) I_{l\pi q}, \\
g_2''' + g_{lq}'''(p_lq) + g_{\pi q}'''(p_\pi q) = e_P''', \\
g_{\pi\pi}'''(p_\pi q) + g_{l\pi}'''(p_lq) = e_P''', \\
g_{ll}'''(p_lq) + g_{l\pi}'''(p_\pi q) = e_l'''.
\end{aligned}$$

Appendix B

In this Appendix we give the formulas for the anti-Hermitian part A_{loop} of the one-loop diagrams. It can be expressed as follows:

$$\begin{aligned}
A_{loop} = \frac{i}{8\pi^2}\sum_n M_{fn}M_{in}^* = \frac{i}{8\pi^2}\sum_n \Bigg[(M_{3a} + M_{3b})(M_{IR} + M_{mag} + M_{Low})\big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} \\
+(M_{5a} + M_{5b} + M_{5c})(M_{IR} + M_{mag} + M_{Low})\big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} \\
+M_7(M_{IR} + M_{mag} + M_{Low})\big|_{\substack{p_l\to k_l,\\ p_\pi\to k_\pi}} + (M_{9a} + M_{9b} + M_{9c} + M_{9d} + M_{9e})M_{10}\Bigg] \\
= (A_{3a-IR} + A_{3a-mag} + A_{3a-Low} + A_{3b-IR} + A_{3b-mag} + A_{3b-Low}) \\
+(A_{5a-IR} + A_{5a-mag} + A_{5a-Low} + A_{5b-IR} + A_{5b-mag} + A_{5b-Low} + A_{5c-IR} + A_{5c-mag} + A_{5c-Low}) \\
+(A_{7-IR} + A_{7-mag} + A_{7-Low}) + (A_{9a-10} + A_{9b-10} + A_{9c-10} + A_{9d-10} + A_{9e-10}), \quad \text{(B.1)}
\end{aligned}$$

where

$$\begin{aligned}
A_{3a-IR} = \frac{i}{8\pi^2}\sum_n M_{3a}M_{IR}\big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} = \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(\hat{p}_K + \hat{p}_\pi)(1+\gamma_5) \\
\times\left[c_0(\hat{P} - m_l)\hat{p}_\pi - \hat{C}_1\hat{p}_\pi + \frac{m_l}{p_lq}(a_0(\hat{P} - m_l) - a_P\hat{P})\right]\frac{\hat{P} - m_l}{2p_lq}\hat{e}^* v_l; \quad \text{(B.2)}
\end{aligned}$$

$$\begin{aligned}A_{3a-mag} = \frac{i}{8\pi^2}\sum_n M_{3a}M_{mag}\Big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} = \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(\hat{p}_K+\hat{p}_\pi)(1+\gamma_5)\\ \times\left[2a_P(P^2+2m_l\hat{P})\right]\frac{\hat{P}-m_l}{(2p_lq)^2}\hat{e}^*v_l;\end{aligned}\tag{B.3}$$

$$\begin{aligned}A_{3a-Low} = \frac{i}{8\pi^2}\sum_n M_{3a}M_{Low}\Big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} = \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(1-\gamma_5)\\ \times\left[2a_0(\hat{P}+2m_l)-2a_P\hat{P}+\hat{C}_1(\hat{P}-m_l)\hat{p}_\pi\right]\frac{\hat{P}-m_l}{2p_lq}\hat{e}^*v_l;\end{aligned}\tag{B.4}$$

$$\begin{aligned}A_{3b-IR} = \frac{i}{8\pi^2}\sum_n M_{3b}M_{IR}\Big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} = \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(\hat{p}_K+\hat{p}_\pi)(1+\gamma_5)\\ \times\frac{1}{2}\Bigg[(\hat{P}-m_l)\hat{e}^*\left(\frac{1}{p_lq}\left((\hat{p}_l-m_l)(b_0\hat{P}-\hat{B}_1)-\hat{B}_1\hat{P}\right)-d_0(\hat{p}_l-m_l)\hat{p}_\pi+\hat{D}_1\hat{p}_\pi\right)\\ -\frac{1}{p_lq}\left(\hat{B}_1\hat{e}^*(\hat{p}_l-m_l)\hat{P}-B_{2\mu\nu}\gamma^\mu\hat{e}^*(\hat{p}_l-m_l)\gamma^\nu-B_{2\mu\nu}\gamma^\mu\hat{e}^*\gamma^\nu\hat{P}\right)\\ +\hat{D}_1\hat{e}^*(\hat{p}_l-m_l)\hat{p}_\pi-D_{2\mu\nu}\gamma^\mu\hat{e}^*\gamma^\nu\hat{p}_\pi\Bigg]v_l;\end{aligned}\tag{B.5}$$

$$\begin{aligned}A_{3b-mag} = \frac{i}{8\pi^2}\sum_n M_{3b}M_{mag}\Big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} = \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(\hat{p}_K+\hat{p}_\pi)(1+\gamma_5)\\ \times\frac{1}{2p_lq}\left[B_{2\mu\nu}\gamma^\mu\hat{p}_l\hat{e}^*\gamma^\nu+4m_lB_{2\mu\nu}\gamma^\mu e^{*\nu}-\hat{B}_1\hat{p}_l\hat{e}^*\hat{P}-4m_l(p_le^*)\hat{B}_1-m_l^2\hat{B}_1\hat{e}^*\right]v_l;\end{aligned}\tag{B.6}$$

$$\begin{aligned}A_{3b-Low} = \frac{i}{8\pi^2}\sum_n M_{3b}M_{Low}\Big|_{\substack{q\to k,\\ p_l\to k_l,\\ e\to e_k}} = \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(1-\gamma_5)\\ \times\frac{1}{2}\Big[D_{2\mu\nu}\gamma^\mu(\hat{P}-m_l)\hat{e}^*\gamma^\nu\hat{p}_\pi-\hat{D}_1(\hat{P}-m_l)\hat{e}^*(\hat{p}_l-m_l)\hat{p}_\pi\\ -2B_{2\mu\nu}\gamma^\mu\hat{e}^*\gamma^\nu+2\hat{B}_1\hat{e}^*\hat{P}+2\hat{p}_l\hat{e}^*\hat{B}_1+8m_lB_{1\mu}e^{*\mu}\\ -2b_0(\hat{p}_l\hat{e}^*\hat{P}+4m_l(p_le^*)+m_l^2\hat{e}^*)\Big]v_l;\end{aligned}\tag{B.7}$$

$$\begin{aligned}A_{5a-IR} = \frac{i}{8\pi^2}\sum_n M_{5a}M_{IR}\Big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} = \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu\Big\{(\hat{p}_K+\hat{P}')\\ \times\left[c_0'(p_lP')-a_0'\frac{p_\pi P'}{p_\pi q}\right]-\left[\hat{C}_1'(p_lP')-a_P'\frac{p_\pi P'}{p_\pi q}\hat{P}'\right]\Big\}(1+\gamma_5)v_l\frac{p_\pi e^*}{p_\pi q};\end{aligned}\tag{B.8}$$

$$\begin{aligned} A_{5a-mag} = \frac{i}{8\pi^2}\sum_n M_{5a}M_{mag}\Big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} &= \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0) \\ &\times \bar{u}_\nu(\hat{p}_K+\hat{P}')(1+\gamma_5)\hat{C}_1'\hat{P}'v_l\frac{p_\pi e^*}{2p_\pi q}; \end{aligned} \tag{B.9}$$

$$\begin{aligned} A_{5a-Low} = \frac{i}{8\pi^2}\sum_n M_{5a}M_{Low}\Big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} &= \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu\hat{P}'(1+\gamma_5)v_l\frac{p_\pi e^*}{p_\pi q} \\ &\times\left[a_0' - a_P'\frac{p_\pi P'}{p_\pi q}\right]; \end{aligned} \tag{B.10}$$

$$\begin{aligned} A_{5b-IR} = \frac{i}{8\pi^2}\sum_n M_{5b}M_{IR}\Big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} &= \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu\Big\{(\hat{p}_K+\hat{P}') \\ \times\Bigg[\frac{p_\pi P'}{p_\pi q}(b_0'(p_\pi e^*) - B_{1\mu}'e^{*\mu}) &- \frac{p_\pi e^*}{p_\pi q}(a_0'-a_P') - (p_l p_\pi)(d_0'(p_\pi e^*) - D_{1\mu}'e^{*\mu})\Bigg] \\ - \Bigg[\frac{p_\pi P'}{p_\pi q}(\hat{B}_1'(p_\pi e^*) - B_{2\mu\nu}'\gamma^\mu e^{*\nu}) &- \frac{1}{p_\pi q}(a_P'(p_\pi e^*)\hat{P}' - A_{2\mu\nu}'\gamma^\mu e^{*\nu}) \\ &-(p_l p_\pi)((p_\pi e^*)\hat{D}_1' - D_{2\mu\nu}'\gamma^\mu e^{*\nu})\Bigg]\Big\}(1+\gamma_5)v_l; \end{aligned} \tag{B.11}$$

$$\begin{aligned} A_{5b-mag} = \frac{i}{8\pi^2}\sum_n M_{5b}M_{mag}\Big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} &= \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(\hat{p}_K+\hat{P}')(1+\gamma_5) \\ &\times\frac{1}{2}\left[D_{2\mu\nu}'\gamma^\mu e^{*\nu} - (p_\pi e^*)\hat{D}_1'\right]\hat{p}_\pi v_l; \end{aligned} \tag{B.12}$$

$$\begin{aligned} A_{5b-Low} = \frac{i}{8\pi^2}\sum_n M_{5b}M_{Low}\Big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} &= \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0) \\ \bar{u}_\nu\Bigg[\frac{p_\pi P'}{p_\pi q}(\hat{B}_1'(p_\pi e^*) - B_{2\mu\nu}'\gamma^\mu e^{*\nu}) &- \frac{1}{p_\pi q}(a_P'(p_\pi e^*)\hat{P}' - A_{2\mu\nu}'\gamma^\mu e^{*\nu}) \\ &-\hat{p}_\pi((p_\pi e^*)b_0' - B_{1\mu}'e^{*\mu})\Bigg](1+\gamma_5)v_l; \end{aligned} \tag{B.13}$$

$$\begin{aligned} A_{5c-IR} = \frac{i}{8\pi^2}\sum_n M_{5c}M_{IR}\Big|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} &= \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu\Big\{(\hat{p}_K+\hat{P}') \\ \times\left[\frac{p_\pi e^*}{p_\pi q}(a_0'-a_P') - (p_l e^*)c_0'\right] &- \left[\frac{p_\pi e^*}{p_\pi q}a_P'\hat{P}' - \frac{1}{p_\pi q}A_{2\mu\nu}'\gamma^\mu e^{*\nu} - (p_l e^*)\hat{C}_1'\right]\Big\}(1+\gamma_5)v_l; \end{aligned} \tag{B.14}$$

$$A_{5c-mag} = \frac{i}{8\pi^2}\sum_n M_{5c}M_{mag}|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} = \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\left[-\frac{1}{2}\bar{u}_\nu(\hat{p}_K+\hat{P}')(1+\gamma_5)\hat{C}_1'\hat{e}^* v_l\right]; \tag{B.15}$$

$$\begin{aligned} A_{5c-Low} = \frac{i}{8\pi^2}\sum_n M_{5c}M_{Low}|_{\substack{q\to k,\\ p_\pi\to k_\pi,\\ e\to e_k}} &= \frac{ie^2}{4\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\\ &\times \bar{u}_\nu\left[\frac{p_\pi e^*}{p_\pi q}a_P'\hat{P}' - \frac{1}{p_\pi q}A'_{2\mu\nu}\gamma^\mu e^{*\nu} - a_0'\hat{e}^*\right](1+\gamma_5)v_l; \end{aligned} \tag{B.16}$$

$$\begin{aligned} A_{7-IR} = \frac{i}{8\pi^2}\sum_n M_7 M_{IR}|_{\substack{p_l\to k_l,\\ p_\pi\to k_\pi}} &= \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(1-\gamma_5)(\hat{p}_K+\hat{P}''+m_l)\\ \times\left[\left(E''_{2\mu\nu}+F''_{2\mu\nu}\right)\gamma^\mu e^{*\nu} - m_l E''_{1\mu}e^{*\mu} - (\hat{P}''-m_l)F''_{1\mu}e^{*\mu}\right]&(\hat{p}_\pi+\hat{P}''+m_l)v_l; \end{aligned} \tag{B.17}$$

$$\begin{aligned} A_{7-mag} = \frac{i}{8\pi^2}\sum_n M_7 M_{mag}|_{\substack{p_l\to k_l,\\ p_\pi\to k_\pi}} &= \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\frac{1}{2}\bar{u}_\nu(1-\gamma_5)\\ \times\left[(\hat{p}_K+\hat{P}'')\hat{q}\hat{e}^*\left(\hat{E}_1''-m_l e_0''\right)+m_l\hat{E}_1''\hat{q}\hat{e}^* - E''_{2\mu\nu}\gamma^\mu\hat{q}\hat{e}^*\gamma^\nu\right]&(\hat{p}_\pi+\hat{P}''+m_l)v_l; \end{aligned} \tag{B.18}$$

$$\begin{aligned} A_{7-Low} = \frac{i}{8\pi^2}\sum_n M_7 M_{Low}|_{\substack{p_l\to k_l,\\ p_\pi\to k_\pi}} &= \frac{ie^2}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c e f_+(0)\bar{u}_\nu(1-\gamma_5)\\ &\times\left[F''_{2\mu\nu}\hat{q}\gamma^\mu e^{*\nu} - \hat{q}(\hat{P}''-m_l)F''_{1\mu}e^{*\mu}+\hat{e}^*\hat{D}_1''\right](\hat{p}_\pi+\hat{P}''+m_l)v_l; \end{aligned} \tag{B.19}$$

$$\begin{aligned} A_{9a-10} = \frac{i}{8\pi^2}\sum_n M_{9a}M_{10} &= \frac{ie^3}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c f_+(0)\bar{u}_\nu(1-\gamma_5)(\hat{p}_K+\hat{P}'''+m_l)\\ &\times\left(\hat{D}_1'''-m_l d_0'''\right)(\hat{p}_\pi+\hat{P}'''+m_l)\left(\frac{p_l e^*}{p_l q}+\frac{\hat{q}\hat{e}^*}{2p_l q}\right)v_l; \end{aligned} \tag{B.20}$$

$$\begin{aligned} A_{9b-10} = \frac{i}{8\pi^2}\sum_n M_{9b}M_{10} &= -\frac{ie^3}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c f_+(0)\bar{u}_\nu(1-\gamma_5)(\hat{p}_K+\hat{P}'''+m_l)\\ \times\Big[\left(F'''_{2\mu\nu}\gamma^\mu e^{*\nu}-m_l F'''_{1\mu}e^{*\mu}\right)&(\hat{p}_\pi+\hat{P}'''+m_l)+\frac{1}{2}\left(\hat{F}_1'''-m_l f_0'''\right)\hat{q}\hat{e}^*(\hat{p}_\pi+\hat{P}''')\\ &+\frac{1}{2}m_l\hat{q}\hat{e}^*\hat{F}_1'''-\frac{1}{2}F'''_{2\mu\nu}\gamma^\mu\hat{q}\hat{e}^*\gamma^\nu\Big]v_l; \end{aligned} \tag{B.21}$$

$$A_{9c-10} = \frac{i}{8\pi^2}\sum_n M_{9c}M_{10} = -\frac{ie^3}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c f_+(0)\bar{u}_\nu(1-\gamma_5)(\hat{p}_K + \hat{P}''' + m_l)$$
$$\times\left((\hat{P}''' - m_l)e_0''' - \hat{E}_1'''\right)(\hat{p}_\pi + \hat{q} + \hat{P}''' + m_l)v_l\frac{p_\pi e^*}{p_\pi q}; \quad \text{(B.22)}$$

$$A_{9d-10} = \frac{i}{8\pi^2}\sum_n M_{9d}M_{10} = \frac{ie^3}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c f_+(0)\bar{u}_\nu(1-\gamma_5)(\hat{p}_K + \hat{P}''' + m_l)$$
$$\times\left[(\hat{P}''' - m_l)G_{1\mu}'''e^{*\mu} - G_{2\mu\nu}'''\gamma^\mu e^{*\nu}\right](\hat{p}_\pi - \hat{q} + \hat{P}''' + m_l)v_l; \quad \text{(B.23)}$$

$$A_{9e-10} = \frac{i}{8\pi^2}\sum_n M_{9e}M_{10} = \frac{ie^3}{8\pi^2}\frac{G}{\sqrt{2}}\sin\theta_c f_+(0)\cdot 2\bar{u}_\nu(1-\gamma_5)(\hat{p}_K + \hat{P}''' + m_l)$$
$$\times\left((\hat{P}''' - m_l)e_0''' - \hat{E}_1'''\right)\hat{e}^* v_l. \quad \text{(B.24)}$$

References

[1] H.W. Fearing, E. Fischbach, and J. Smith, Phys. Rev. **D 2**, 542 (1970).

[2] H.W. Fearing, E. Fischbach, and J. Smith, Phys. Rev. Lett. **24**, 189 (1970).

[3] J. Bijnens, G. Ecker, and J. Gasser, Nucl. Phys. **B 396**, 81 (1993).

[4] J. Gasser, B. Kubis, N. Paver, and M. Verbeni, Eur. Phys. J. **C 40**, 205 (2005).

[5] V.V. Braguta, A.A. Likhoded, and A.E. Chalov, Phys. Rev. **D 65**, 054038 (2002).

[6] E.H. Müller, B. Kubis, and U.-G. Meißner, Eur. Phys. J. **C 48**, 427 (2006).

[7] I.B. Khriplovich and A.S. Rudenko, arXiv:1012.0147 [hep-ph], to be published in Phys. Atom. Nucl.

[8] F.E. Low, Phys. Rev. **110**, 974 (1958).

[9] K. Nakamura *et al.* (Particle Data Group), J. Phys. **G 37**, 075021 (2010).

[10] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. **B 592**, 634 (2004).

[11] S.A. Akimenko *et al.*, Phys. Atom. Nucl. **70**, 702 (2007).

[12] O.G. Tchikilev *et al.*, Phys. Atom. Nucl. **70**, 29 (2007).